

University of Groningen

Atorvastatin (Lipitor) by MCR

Zarganes-Tzitzikas, Tryfon; Neochoritis, Constantinos G.; Dömling, Alexander

Published in:
Bioorganic & Medicinal Chemistry Letters

DOI:
[10.1021/acsmedchemlett.8b00579](https://doi.org/10.1021/acsmedchemlett.8b00579)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Zarganes-Tzitzikas, T., Neochoritis, C. G., & Dömling, A. (2019). Atorvastatin (Lipitor) by MCR. *Bioorganic & Medicinal Chemistry Letters*, 10(3), 389-392. <https://doi.org/10.1021/acsmedchemlett.8b00579>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


Atorvastatin (Lipitor) by MCR

Tryfon Zarganes-Tzitzikas, Constantinos G. Neochoritis,¹ and Alexander Dömling^{*,2}

Department of Pharmacy, Drug Design group, University of Groningen, A. Deusinglaan 1, Groningen 9700 AV, The Netherlands

S Supporting Information

ABSTRACT: A concise and convergent synthesis of the atorvastatin, the best-selling cardiovascular drug of all time, is presented. Our approach is based on an Ugi reaction, which shortens the current synthetic route and is advantageous over the published syntheses.



KEYWORDS: Atorvastatin, Ugi reaction, münchenone, convergent synthesis, generics

Multicomponent reactions (MCRs) are an advanced class of organic reactions which, opposite to classical organic reactions, allow for the easy, fast, and efficient generation of chemical diversity in just one assembly step.^{1–3} These features make them an attractive area in research and development.⁴ Surprisingly, the number of applications in drug discovery is rather limited regarding the superb advantages of this chemistry.⁵ An analysis of the currently marketed drugs, however, shows that approximately 5% can be synthesized with the use of MCR, even so they are synthesized by a classical sequential pathway.⁶ Examples of drugs synthesized by MCR clearly show the immense advantages of them in this context, e.g., lidocaine,⁷ praziquantel,^{8,9} telaprevir,¹⁰ olanzapine,¹¹ clopidogrel,¹² lacosamide,¹³ carfentanil,¹⁴ ivosidenib,¹⁵ and levetiracetam (Figure 1).¹⁶ Epelsiban¹⁷ and almorexant¹⁸ are examples of compounds currently or recently in clinical trials and actually synthesized by utilization of the MCR repertoire (Figure 1).

Here we report an MCR-based synthesis of atorvastatin (common trade name: Lipitor), one of the world's best-selling medication of all time. Only in 2005, Lipitor made \$12 billion in sales and was used by more than 45 million people worldwide.¹⁹ It belongs to the drug class of statins, lipid-lowering drugs for the prevention of events associated with cardiovascular disease.²⁰ It is an example of a competitive HMG-CoA-reductase inhibitor, which consists of a pentasubstituted pyrrole core. The importance of atorvastatin until today^{21–23} led to much interest in its synthesis. The main retrosynthetic scheme of the atorvastatin synthesis as described in literature focuses on the assembly of its five different substituents on a pyrrole hub.^{24,25} By this way, which consists also the industrial route,²⁶ the pyrrole ring could be formed by a Paal–Knorr cyclocondensation^{27,28} of the highly substituted 1,4-diketone **2** with primary amine **3** (Paal–Knorr route, Scheme 1, blue color).^{21,22,26,29–34} In 2015, a total synthesis of atorvastatin via a late-stage, regioselective 1,3-dipolar münchenone cycloaddition³⁵ of the amido acid **4** with the acetylene




Figure 1. Examples of marketed drugs and drugs in clinical trials which have been discovered using MCR chemistry; the amine, aldehyde, isocyanide, and acid components are depicted with green, red, blue, and magenta color, respectively.

derivative **5** (münchenone route, Scheme 1, red color) was described.³⁶ Although this synthesis provided a nice solution to the problem of regioselectivity of the cycloaddition,³⁰ the synthesis of derivative **4** required five sequential steps which contributed to eight steps for the total synthesis of atorvastatin. Regarding the latter approach, we envisioned the synthesis of the amido acid **4** in only two steps utilizing the Ugi four-component reaction (U-4C, Scheme 2).


The initial derived MCR adduct can be considered as a synthetic hub to a vast diversity of other scaffolds.² Thus, the 1,4 amido acid motif could easily be derived from an Ugi adduct with the cleavage of the isocyanide moiety (Scheme 2).^{37,38} Indeed, the reaction at rt of *p*-fluorobenzaldehyde **6**,

Received: November 26, 2018


Accepted: February 7, 2019

Published: February 7, 2019

Scheme 1. Main Retrosynthetic Scheme for the Synthesis of Atorvastatin (Paal–Knorr Route, Blue Color); A Novel Approach to the Intermediate 4 Is Proposed by MCR (Münchnone Route, Red Color)



Scheme 2. MCR-Based Synthesis of 4 and the Subsequent Synthesis Towards Atorvastatin 1



the suitably functionalized, commercially available amine **3**,^{29,39–41} the convertible isocyanide **7**,^{42–45} and isobutyric acid **8** in 2,2,2-trifluoroethanol (TFE) afforded the Ugi adduct (U-4C) **9** in 40% yield. The choice of the corresponding isocyanide was the easiness of its cleavage at basic pH, keeping intact the other functional groups. Thus, in a one-pot acid deprotection and isocyanide cleavage, we obtained the valuable intermediate **4** in a *dr* 5:4 in 87% yield. Then, we performed the regioselective [3 + 2] cycloaddition³⁶ of **4** with the *N*,3-diphenylpropiolamide **5** and *N*,*N*'-diisopropylcarbodiimide (DIPC) in THF, yielding the advanced intermediate **10** in 46% yield which can be readily converted by acidic deprotection with 10-camphorsulfonic acid (CSA) to atorvastatin **1** (Scheme 2).

The industrial atorvastatin synthesis via the Paal–Knorr route is a synthesis consisting of six steps excluding the synthesis of amine **3**, which is commercially available (Table 1). MCR chemistry has also been employed in order to improve and optimize this synthetic route. These modifications include a one-pot Stetter/Paal–Knorr reaction sequence catalyzed by NHC⁴⁶ or a Hantzsch pyrrole synthesis (Table

1).⁴⁷ Regarding the münchnone route, this is the first time to the best of our knowledge, that MCR chemistry is utilized. On the basis of MCR chemistry, we synthesized the intermediate **4** in only two steps, and with two additional steps, we successfully obtained atorvastatin (Scheme 2). The Ugi reaction was performed at 10 mmol scale, see Supporting Information).

Our current approach effectively reduces the number of steps toward atorvastatin to only four compared with the seven reported in literature and establish this methodology equally or even better than the Paal–Knorr route. We can classify the recent syntheses of atorvastatin in four different routes (Table 1). Most of the published Paal–Knorr route syntheses include different variations of the synthesis of the amine (entry 1) or differentiation in the amine vector of the pyrrole core (entries 1–3). The required steps vary from six to 10. A Stetter/Paal–Knorr reaction sequence (entry 4) and a Hantzsch pyrrole synthesis (entry 5) were presented as alternatives with four and five steps, respectively. Our synthetic strategy can be ranked among the most competitive one with four steps (entry 7).⁴⁸

Table 1. Comparison of the Most Important, Recent Atorvastatin Syntheses in Literature along with Our MCR Approach

	routes	reference/ report	steps	remarks
1	Paal–Knorr	22,34,29 ^a	6 ^b	different variations on the synthesis of amine 3/ differentiation in the amine vector of the pyrrole core
2		40	8	differentiation in the amine vector of the pyrrole core
3		23	10	differentiation in the amine vector of the pyrrole core
4	Stetter/ Paal–Knorr	46	4 ^{b,c}	NHC-catalyzed Stetter/ Paal–Knorr sequence
5	Hantzsch	47	5 ^d	Hantzsch variation of the pyrrole synthesis
6	Münchnone	36	7	
7		this work	4	

^aThe corresponding methyl ester of the amine 3 was employed in the Paal–Knorr ^bExcluding the steps required for the synthesis of amine 3

^cThe final product of the synthesis is the fully protected atorvastatin.

^dThe final product of the synthesis is the atorvastatin lactone.

It is noteworthy that our current synthetic methodology of utilizing an MCR adduct bears convertible isocyanides, yielding the 1,4-amido acid motif. This strategy is beneficial not only because we have a faster access to atorvastatin but also by this way more derivatives are accessible. Thus, we can readily synthesize substituted bioactive pyrroles with a great diversity on substituents in 1-, 2-, and 5-positions, for example, positron emission tomography (PET) labeled derivatives.³⁶

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsmchemlett.8b00579.

Experimental procedures and full characterization for compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Phone: +31-6-3633307. E-mail: a.s.s.domling@rug.nl.

ORCID

Constantinos G. Neochoritis: 0000-0001-5098-5504

Alexander Dömling: 0000-0002-9923-8873

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding

This research has been supported to (AD) by the National Institute of Health (NIH) (2R01GM097082-05), the European Lead Factory (IMI) under grant agreement number 115489, the Qatar National Research Foundation (NPRP6-065-3-012). Moreover funding was received through ITN “Accelerated Early stage drug discovery” (AEGIS, grant agreement no. 675555) and COFUNDs ALERT and PROMINENT (grant agreements no. 665250 and 754425), Hartstichting (ESCAPE-HF, 2018B012) and KWF Kankerbestrijding grant (grant agreement no. 10504).

Notes

The authors declare no competing financial interest.

■ ABBREVIATIONS USED

TFE, 2,2,2-trifluoroethanol; DIPIC, *N,N'*-diisopropylcarbodiimide; CSA, 10-camphorsulfonic acid; DCM, dichloromethane; PET, positron emission tomography.

■ REFERENCES

- (1) Dömling, A.; Wang, W.; Wang, K. Chemistry and Biology Of Multicomponent Reactions. *Chem. Rev.* **2012**, *112*, 3083–3135.
- (2) Dömling, A. Recent Developments in Isocyanide Based Multicomponent Reactions in Applied Chemistry. *Chem. Rev.* **2006**, *106*, 17–89.
- (3) Dömling, A.; Ugi, I. Multicomponent Reactions with Isocyanides. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168–3210.
- (4) Cioc, R. C.; Ruijter, E.; Orru, R. V. A. Multicomponent Reactions: Advanced Tools for Sustainable Organic Synthesis. *Green Chem.* **2014**, *16*, 2958–2975.
- (5) Brown, D. G.; Boström, J. Analysis of Past and Present Synthetic Methodologies on Medicinal Chemistry: Where Have All the New Reactions Gone? *J. Med. Chem.* **2016**, *59*, 4443–4458.
- (6) Zarganes-Tzitzikas, T.; Dömling, A. Modern Multicomponent Reactions for Better Drug Syntheses. *Org. Chem. Front.* **2014**, *1*, 834–837.
- (7) Ugi, I.; Steinbrückner, C. Über Ein Neues Kondensations-Prinzip. *Angew. Chem.* **1960**, *72*, 267–268.
- (8) Dömling, A.; Khoury, K. Praziquantel and Schistosomiasis. *ChemMedChem* **2010**, *5*, 1420–1434.
- (9) Cao, H.; Liu, H.; Dömling, A. Efficient Multicomponent Reaction Synthesis of the Schistosomiasis Drug Praziquantel. *Chem. - Eur. J.* **2010**, *16*, 12296–12298.
- (10) Znabet, A.; Polak, M. M.; Janssen, E.; de Kanter, F. J. J.; Turner, N. J.; Orru, R. V. A.; Ruijter, E. A Highly Efficient Synthesis of Telaprevir by Strategic Use of Biocatalysis and Multicomponent Reactions. *Chem. Commun.* **2010**, *46*, 7918–7920.
- (11) Gewald, K.; Schinke, E.; Böttcher, H. Heterocyclen Aus CH-Aciden Nitrilen, VIII. 2-Amino-Thiophene Aus Methylenaktiven Nitrilen, Carbonylverbindungen Und Schwefel. *Chem. Ber.* **1966**, *99*, 94–100.
- (12) Gurbel, P. A.; O'Connor, C. M.; Cummings, C. C.; Serebruany, V. L. Clopidogrel: The Future Choice for Preventing Platelet Activation during Coronary Stenting? *Pharmacol. Res.* **1999**, *40*, 107–111.
- (13) Wehlan, H.; Oehme, J.; Schäfer, A.; Rossen, K. Development of Scalable Conditions for the Ugi Reaction—Application to the Synthesis of (R)-Lacosamide. *Org. Process Res. Dev.* **2015**, *19*, 1980–1986.
- (14) Váradi, A.; Palmer, T. C.; Haselton, N.; Afonin, D.; Subrath, J. J.; Le Rouzic, V.; Hunkele, A.; Pasternak, G. W.; Marrone, G. F.; Borics, A.; et al. Synthesis of Carfentanil Amide Opioids Using the Ugi Multicomponent Reaction. *ACS Chem. Neurosci.* **2015**, *6*, 1570–1577.
- (15) Popovici-Muller, J.; Lemieux, R. M.; Artin, E.; Saunders, J. O.; Salituro, F. G.; Travins, J.; Cianchetta, G.; Cai, Z.; Zhou, D.; Cui, D.; et al. Discovery of AG-120 (Ivosidenib): A First-in-Class Mutant IDH1 Inhibitor for the Treatment of IDH1 Mutant Cancers. *ACS Med. Chem. Lett.* **2018**, *9*, 300–305.
- (16) Cioc, R.; Schaepkens van Riepmst, L.; Schuckman, P.; Ruijter, E.; Orru, R. Ugi Four-Center Three-Component Reaction as a Direct Approach to Racetams. *Synthesis* **2017**, *49*, 1664–1674.
- (17) Borthwick, A. D.; Liddle, J. Retosiban and Epelsiban: Potent and Selective Orally Available Oxytocin Antagonists. In *Protein-Protein Interactions in Drug Discovery*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2013; pp 225–256.
- (18) Aissaoui, H.; Cappi, M.; Clozel, M.; Fischli, W.; Ralf, K. 1,2,3,4-Tetrahydroisoquinoline Derivatives. World Patent WO2001068609A120010920, 2000.

- (19) Simons, J. The \$10 Billion Pill Hold the Fries, Please. Lipitor, the Cholesterol-Lowering Drug, Has Become the Bestselling Pharmaceutical in History. Here's How Pfizer Did It. *Fortune* January 20, 2003.
- (20) Collins, R.; Reith, C.; Emberson, J.; Armitage, J.; Baigent, C.; Blackwell, L.; Blumenthal, R.; Danesh, J.; Smith, G. D.; DeMets, D.; et al. Interpretation of the Evidence for the Efficacy and Safety of Statin Therapy. *Lancet* **2016**, 388, 2532–2561.
- (21) Novozhilov, Y. V.; Dorogov, M. V.; Blumina, M. V.; Smirnov, A. V.; Krasavin, M. An Improved Kilogram-Scale Preparation of Atorvastatin Calcium. *Chem. Cent. J.* **2015**, 9, 7.
- (22) Li, F.; Jiang, W.; Czarnik, A. W.; Li, W. Combinatorial Synthesis of Deuterium-Enriched Atorvastatin. *Mol. Diversity* **2016**, 20, 453–459.
- (23) Dias, L. C.; Vieira, A. S.; Barreiro, E. J. The Total Synthesis of Calcium Atorvastatin. *Org. Biomol. Chem.* **2016**, 14, 2291–2296.
- (24) Baumann, M.; Baxendale, I. R.; Ley, S. V.; Nikbin, N. An Overview of the Key Routes to the Best Selling 5-Membered Ring Heterocyclic Pharmaceuticals. *Beilstein J. Org. Chem.* **2011**, 7, 442–495.
- (25) Roth, B. D. The Discovery and Development of Atorvastatin, A Potent Novel Hypolipidemic Agent. In *Progress in Medicinal Chemistry*; Elsevier, **2002**; Vol. 40, pp 1–22, DOI: .
- (26) Butler, D. E.; Le, T. V.; Millar, A.; Nanninga, T. N. Process for the Synthesis of (5R)-1,1-Dimethylethyl-6-cyano-5-hydroxy-3-oxo-hexanoate. U.S. Patent US5155251A, 1992, no. 19.
- (27) Knorr, L. Synthese von Furfuranderivaten Aus Dem Diacetbernsteinsäureester. *Ber. Dtsch. Chem. Ges.* **1884**, 17, 2863–2870.
- (28) Paal, C. Ueber Die Derivate Des Acetophenonacetessigesters Und Des Acetonylacetessigesters. *Ber. Dtsch. Chem. Ges.* **1884**, 17, 2756–2767.
- (29) Kawato, Y.; Iwata, M.; Yazaki, R.; Kumagai, N.; Shibasaki, M. A Simplified Catalytic System for Direct Catalytic Asymmetric Aldol Reaction of Thioamides; Application to an Enantioselective Synthesis of Atorvastatin. *Tetrahedron* **2011**, 67, 6539–6546.
- (30) Roth, B. D.; Blankley, C. J.; Chucholowski, A. W.; Ferguson, E.; Hoefle, M. L.; Ortwine, D. F.; Newton, R. S.; Sekerke, C. S.; Sliskovic, D. R.; Wilson, M. Inhibitors of Cholesterol Biosynthesis. 3. Tetrahydro-4-Hydroxy-6-[2-(1H-Pyrrol-1-Yl)Ethyl]-2H-Pyran 2-One Inhibitors of HMG-CoA Reductase. 2. Effects of Introducing Substituents at Positions Three and Four of the Pyrrole Nucleus. *J. Med. Chem.* **1991**, 34, 357–366.
- (31) Brower, P. L.; Butler, D. E.; Deering, C. F.; Le, T. V.; Millar, A.; Nanninga, T. N.; Roth, B. D. The Synthesis of (4R-Cis)-1,1-Dimethylethyl 6-Cyanomethyl-2,2-Dimethyl-1,3-Dioxane-4-Acetate, a Key Intermediate for the Preparation of CI-981, a Highly Potent, Tissue Selective Inhibitor of HMG-CoA Reductase. *Tetrahedron Lett.* **1992**, 33, 2279–2282.
- (32) Baumann, K. L.; Butler, D. E.; Deering, C. F.; Mennen, K. E.; Millar, A.; Nanninga, T. N.; Palmer, C. W.; Roth, B. D. The Convergent Synthesis of CI-981, an Optically Active, Highly Potent, Tissue Selective Inhibitor of HMG-CoA Reductase. *Tetrahedron Lett.* **1992**, 33, 2283–2284.
- (33) Roth, B. D. Trans-6-[2-(3- or 4-Carboxamido-substituted pyrrol-1-yl)alkyl]-4-hydroxypyran-2-one Inhibitors of Cholesterol Synthesis. U.S. Patent US 4,681,893 C1, 1987.
- (34) Sagyam, R. R.; Padi, P. R.; Ghanta, M. R.; Vurimidi, H. An Efficient Synthesis of Highly Substituted Pyrrole and Bis Pyrrole Derivatives. *J. Heterocycl. Chem.* **2007**, 44, 923–926.
- (35) Pandey, P.; Srinivasa Rao, T. An Efficient Synthesis of N3,4-Diphenyl-5-(4-Fluorophenyl)-2-Isopropyl-1H-3-Pyrrolicarboxamide, a Key Intermediate for Atorvastatin Synthesis. *Bioorg. Med. Chem. Lett.* **2004**, 14, 129–131.
- (36) Lopchuk, J. M.; Gribble, G. W. Total Synthesis of Atorvastatin via a Late-Stage, Regioselective 1,3-Dipolar Münchnone Cycloaddition. *Tetrahedron Lett.* **2015**, 56, 3208–3211.
- (37) Keating, T. A.; Armstrong, R. W. Postcondensation Modifications of Ugi Four-Component Condensation Products: 1-Isocyanocyclohexene as a Convertible Isocyanide. Mechanism of Conversion, Synthesis of Diverse Structures, and Demonstration of Resin Capture. *J. Am. Chem. Soc.* **1996**, 118, 2574–2583.
- (38) van der Heijden, G.; Sjaak Jong, J. A. W.; Ruijter, E.; Orru, R. V. A. 2-Bromo-6-Isocyanopyridine as a Universal Convertible Isocyanide for Multicomponent Chemistry. *Org. Lett.* **2016**, 18, 984–987.
- (39) Chen, X.; Xiong, F.; Chen, W.; He, Q.; Chen, F. Asymmetric Synthesis of the HMG-CoA Reductase Inhibitor Atorvastatin Calcium: An Organocatalytic Anhydride Desymmetrization and Cyanide-Free Side Chain Elongation Approach. *J. Org. Chem.* **2014**, 79, 2723–2728.
- (40) Gao, J.; Guo, Y. H.; Wang, Y. P.; Wang, X. J.; Xiang, W. S. A Novel and Efficient Route for the Preparation of Atorvastatin. *Chin. Chem. Lett.* **2011**, 22, 1159–1162.
- (41) Müller, M. Chemoenzymatic Synthesis of Building Blocks for Statin Side Chains. *Angew. Chem., Int. Ed.* **2005**, 44, 362–365.
- (42) Baldoli, C.; Giannini, C.; Licandro, E.; Maiorana, S.; Zinzalla, G. A Thymine-PNA Monomer as New Isocyanide Component in the Ugi Reaction: A Direct Entry to PNA Dimers. *Synlett* **2004**, 2004, 1044–1048.
- (43) Baldoli, C.; Maiorana, S.; Licandro, E.; Zinzalla, G.; Perdicchia, D. Synthesis of Chiral Chromium Tricarbonyl Labeled Thymine PNA Monomers via the Ugi Reaction. *Org. Lett.* **2002**, 4, 4341–4344.
- (44) Maison, W.; Schlemminger, I.; Westerhoff, O.; Martens, J. Multicomponent Synthesis of Novel Amino Acid–nucleobase Chimeras: A Versatile Approach to PNA-Monomers. *Bioorg. Med. Chem.* **2000**, 8, 1343–1360.
- (45) Maison, W.; Schlemminger, I.; Westerhoff, O.; Martens, J. Modified PNAs: A Simple Method for the Synthesis of Monomeric Building Blocks. *Bioorg. Med. Chem. Lett.* **1999**, 9, 581–584.
- (46) Fleige, M.; Glorius, F. α -Unsubstituted Pyrroles by NHC-Catalyzed Three-Component Coupling: Direct Synthesis of a Versatile Atorvastatin Derivative. *Chem. - Eur. J.* **2017**, 23, 10773–10776.
- (47) Estévez, V.; Villacampa, M.; Menéndez, J. C. Concise Synthesis of Atorvastatin Lactone under High-Speed Vibration Milling Conditions. *Org. Chem. Front.* **2014**, 1, 458–463.
- (48) Dömling, A. Methods for Providing Intermediates in the Synthesis of Atorvastatin. World Patent WO2016122325A1, 2016.